

PRIOR AUTHORIZATION POLICY

- POLICY:** Immunologicals – Ebglyss Prior Authorization Policy
- Ebglyss® (lebrikizumab-lbkz subcutaneous injection – Eli Lilly)

REVIEW DATE: 09/18/2024

OVERVIEW

Ebglyss, an interleukin (IL)-13 antagonist, is indicated for the treatment of moderate to severe atopic dermatitis in adults and pediatric patients ≥ 12 years of age who weigh ≥ 40 kg whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable.¹ Ebglyss may be used with or without topical corticosteroids (TCSs).

Clinical Efficacy

Three pivotal studies of Ebglyss enrolled patients ≥ 12 years of age with moderate to severe chronic atopic dermatitis affecting $\geq 10\%$ of their body surface area.¹⁻⁴ Patients also had a of an inadequate response to a sufficient course of topical therapy (e.g., topical corticosteroids). At Week 16, Ebglyss was found to be more effective in achieving a clinical response compared with placebo. In the monotherapy trials, the majority of patients who achieved a clinical response to Ebglyss at Week 16 experienced sustained efficacy at Week 52.

Guidelines

Current atopic dermatitis guidelines do not make recommendations regarding Ebglyss. The **American Academy of Dermatology (AAD) Guidelines for the Care and Management of Atopic Dermatitis in Adults (2023)** and the **American Academy of Allergy, Asthma and Immunology (AAAAI)/American College of Allergy, Asthma and Immunology (ACAAI) Joint Task Force on Practice Parameters Atopic Dermatitis Guidelines (2023)** continue to affirm that despite the availability of newer, systemic therapies, topical agents remain the mainstay of treatment due to their proven track record and favorable safety profiles.⁵⁻⁷ Several topical agents are recommended, with topical corticosteroids commonly used first-line for mild to severe atopic dermatitis in all skin regions. If topical therapy and basic management (e.g., moisturizers, bathing modifications) have been optimized and the patient has not achieved adequate control, systemic therapy may be considered.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Ebglyss. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Ebglyss as well as the monitoring required for adverse events and long-term efficacy, initial approval requires Ebglyss to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

09/18/2024

© 2024. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Ebglyss is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Atopic Dermatitis.** Approve for the duration noted if the patient meets ONE of the following conditions (A or B):
 - A) **Initial Therapy.** Approve for 4 months if the patient meets ALL of the following criteria (i, ii, iii, and iv):
 - i. Patient meets ONE of the following criteria (a or b):
 - a) Patient is ≥ 18 years of age; OR
 - b) Patient meets BOTH of the following criteria (1 and 2):
 - 1) Patient is 12 to 17 years of age; AND
 - 2) Patient weighs ≥ 40 kg; AND
 - ii. Patient has atopic dermatitis involvement estimated to be $\geq 10\%$ of the body surface area according to the prescriber; AND
 - iii. Patient meets ALL of the following criteria (a, b, and c):
 - a) Patient has tried at least one medium-, medium-high, high-, and/or super-high-potency prescription topical corticosteroid; AND
 - b) This topical corticosteroid was applied daily for at least 28 consecutive days; AND
 - c) Inadequate efficacy was demonstrated with this topical corticosteroid therapy, according to the prescriber; AND
 - iv. The medication is prescribed by or in consultation with an allergist, immunologist, or dermatologist.
 - B) **Patient is Currently Receiving Ebglyss.** Approve for 1 year if the patient meets BOTH of the following criteria (i and ii):
 - i. Patient has already received at least 4 months of therapy with Ebglyss; AND
Note: A patient who has received < 4 months of therapy or who is restarting therapy with Ebglyss should be considered under criterion 1A (Atopic Dermatitis, Initial Therapy).
 - ii. Patient has responded to therapy as determined by the prescriber.
Note: Examples of a response to Ebglyss therapy are marked improvements in erythema, induration/papulation/edema, excoriations, and lichenification; reduced pruritus; decreased requirement for other topical or systemic therapies; reduced body surface area affected with atopic dermatitis; or other responses observed.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Ebglyss is not recommended in the following situations:

1. **Asthma.** Ebglyss is not indicated for the treatment of asthma.¹ Several studies, including one Phase III study, evaluated Ebglyss for the treatment of adults with asthma with mixed results.⁸⁻¹¹ Two replicate Phase IIb studies (designed as Phase III, but host-cell impurity in the study drug material was discovered), LUTE and VERSE (published) [n = 463] evaluated Ebglyss in patients with uncontrolled asthma despite treatment with medium- to high-dose inhaled corticosteroids and a second asthma controller.¹⁰ Following a mean 24 weeks of treatment, Ebglyss significantly reduced asthma exacerbations vs. placebo in patients with in patients who had baseline periostin levels > 50 ng/mL. However, in the Phase III study, STRETTO (published) [n = 310], Ebglyss did not significantly improve forced expiratory volume in 1 second compared with placebo in patients with mild to moderate asthma.¹¹

2. **Concurrent use of Ebglyss with another Monoclonal Antibody Therapy (i.e., Adbry, Dupixent, Cinqair, Fasenra, Nemluvio, Nucala, Tezspire, or Xolair).** The efficacy and safety of Ebglyss in combination with other monoclonal antibodies have not been established.
3. **Concurrent Use of Ebglyss with Janus Kinase Inhibitors (JAKis) [oral or topical].** Use of JAK inhibitors is not recommended for use in combination with other JAK inhibitors, biologic immunomodulators (e.g., Ebglyss), or with other immunosuppressants.¹²⁻¹⁵
Note: Examples of JAK inhibitors are Cibinco[®] (abrocitinib tablets), Leqselvi[™] (deuruxolitinib tablets), Rinvoq[®]/Rinvoq[®] LQ (upadacitinib tablets and oral solution), and Opzelura[®] (ruxolitinib cream).
4. **Idiopathic Pulmonary Fibrosis.** Ebglyss is not indicated for the treatment of idiopathic pulmonary fibrosis.¹ In one Phase II, randomized, placebo-controlled study (published) [n = 505], Ebglyss was not found to provide a benefit in forced vital capacity decline vs. placebo when either administered as monotherapy or in combination with pirfenidone.¹⁶
5. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Ebglyss[®] subcutaneous injection [prescribing information]. Indianapolis, IN: Eli Lilly; September 2024.
2. Silverberg JI, Guttman-Yassky E, Thaci D, et al. Two phase 3 trials of lebrikizumab for moderate-to-severe atopic dermatitis. *N Engl J Med.* 2023;388(12):1080-1091.
3. Simpson EL, Gooderham M, Wollenberg A, et al. Efficacy and safety of lebrikizumab in combination with topical corticosteroids in adolescents and adults with moderate to severe atopic dermatitis. *JAMA Dermatol.* 2023;159(2):182-191.
4. Blauvelt A, Thyssen JP, Guttman-Yassky E, et al. Efficacy and safety of lebrikizumab in moderate-to-severe atopic dermatitis: 52-week results of two randomized double-blinded placebo-controlled phase III trials. *B J Dermatol.* 2023;188(6):740-748.
5. Sidbury R, Alikhan A, Cohen DE, et al. Guidelines of care for the management of atopic dermatitis in adults with topical therapies. *J Am Acad Dermatol.* 2023;89(1):e1-e20.
6. Davis DMR, Drucker AM, Alikhan A, et al. Guidelines of care for the management of atopic dermatitis in adults with phototherapy and systemic therapies. *J Am Acad Dermatol.* 2024;90(2):e43-e56.
7. Chu DK, Schneider L, Asiniwasis RN, et al. Atopic dermatitis (eczema) guidelines: 2023 American Academy of Allergy Asthma and Immunology/American College of Allergy, Asthma and Immunology Joint Task Force on practice parameters GRADE- and Institute of Medicine-based recommendations. *Ann Allergy Asthma Immunol.* 2024;132(3):274-312.
8. Corren J, Lemanske RF, Hanania NA, et al. Lebrikizumab treatment in adults with asthma. *N Engl J Med.* 2011;365(12):1088-1098.
9. Noonan M, Korenblat P, Mosesova S, et al. Dose-ranging study of lebrikizumab in asthmatic patients not receiving inhaled steroids. *J Allergy Clin Immunol.* 2013;132(3):567-574.
10. Hanania NA, Noonan M, Corren J, et al. Lebrikizumab in moderate-to-severe asthma: pooled data from two randomized placebo-controlled studies. *Thorax.* 2015;70(8):748-756.
11. Korenblat P, Kerwin E, Leshchenko I, et al. Efficacy and safety of lebrikizumab in adult patients with mild-to-moderate asthma not receiving inhaled corticosteroids. *Respir Med.* 2018;134:143-149.
12. Cibinco[®] tablets [prescribing information]. New York, NY: Pfizer; December 2023.
13. Rinvoq[®] tablets [prescribing information]. North Chicago, IL: AbbVie; April 2024.
14. Opzelura[®] cream [prescribing information]. Wilmington, DE: Incyte; March 2023.
15. Leqselvi[™] tablets [prescribing information]. Whippany, NJ: Sun/Halo; July 2024.
16. Maher TM, Costabel U, Glassberg MK, et al. Phase 2 trial to assess lebrikizumab in patients with idiopathic pulmonary fibrosis. *Eur Respir J.* 2021;57(2):1902442.