

PRIOR AUTHORIZATION POLICY

- POLICY:** Iron Replacement – Ferrlecit Prior Authorization Policy
- Ferrlecit[®] (sodium ferric gluconate complex in sucrose intravenous infusion – sanofi-aventis, generic)

REVIEW DATE: 01/10/2024

OVERVIEW

Ferrlecit, an iron replacement product, is indicated for the treatment of **iron deficiency anemia** in patients ≥ 6 years of age with **chronic kidney disease (CKD) receiving hemodialysis** who are receiving supplemental epoetin therapy.¹

Guidelines

The Kidney Disease: Improving Global Outcomes guidelines for anemia in CKD (2012) make various recommendations regarding iron therapy.² For adults with CKD and anemia not on iron or erythropoietin stimulating agent (ESA) therapy, a trial of intravenous (IV) iron (or in non-dialysis patients with CKD, alternatively, a 1 to 3 month trial of oral iron therapy) is recommended if an increase in hemoglobin (Hb) concentration without starting ESA treatment is desired and transferrin saturation (TSAT) is $\leq 30\%$ and ferritin is ≤ 500 ng/mL. For adults with CKD on ESA therapy who are not receiving iron supplementation, a trial of IV iron (or in non-dialysis CKD patients, alternatively, a 1 to 3 month trial of oral iron therapy) is recommended if an increase in Hb concentration or a decrease in ESA dose is desired and TSAT is $\leq 30\%$ and ferritin is ≤ 500 ng/mL. For all pediatric patients with CKD with anemia not on iron or ESA therapy, oral iron (or IV iron in patients receiving hemodialysis) is recommended when TSAT is $\leq 20\%$ and ferritin is ≤ 100 ng/mL. For all pediatric patients with CKD who are receiving ESA therapy but not receiving iron supplementation, it is recommended to administer oral iron (or IV iron for patients receiving hemodialysis) to maintain TSAT $> 20\%$ and ferritin > 100 ng/dL.

The National Comprehensive Cancer Network guidelines on Hematopoietic Growth Factors (version 12.2024 – December 12, 2023) discuss the management of cancer- and chemotherapy-induced anemia.³ Treatment for iron deficiency is guided by iron status which is defined in the guidelines as: absolute iron deficiency, functional iron deficiency, possible functional iron deficiency, or no iron deficiency. IV iron therapy is considered an option for patients with absolute iron deficiency (ferritin < 30 ng/mL and TSAT $< 20\%$) in patients who are also receiving an ESA, functional iron deficiency (ferritin = 30 to 500 ng/mL and TSAT $< 50\%$), and for select patients with possible functional iron deficiency (ferritin = 501 to 800 ng/mL and TSAT $< 50\%$).

The American College of Cardiology/American Heart Association guideline for the management of heart failure (2022) states that in patients with heart failure with reduced ejection fraction (left ventricular ejection fraction $\leq 40\%$), absolute iron deficiency (ferritin < 100 ng/mL) or functional iron deficiency (ferritin = 100 to 300 mg/mL if TSAT is $< 20\%$), and with or without anemia, IV iron replacement is reasonable to improve functional status and quality of life (2a recommendation).⁴

01/10/2024

© 2024. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Ferrlecit. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Ferrlecit as well as the monitoring required for adverse events and long-term efficacy, particular approvals require Ferrlecit to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Ferrlecit is recommended in those who meet one of the following criteria:

FDA-Approved Indication

- 1. Iron Deficiency Anemia in Patients with Chronic Kidney Disease who are on Dialysis.** Approve for 3 years.

Other Uses with Supportive Evidence

- 2. Iron Deficiency Anemia in Patients with Chronic Kidney Disease who are not on Dialysis.** Approve for 1 year if the patient meets the following (A and B):
 - A)** Patient is ≥ 6 years of age; AND
 - B)** Ferrlecit is prescribed by or in consultation with a nephrologist or hematologist.
- 3. Iron Deficiency Anemia, Other.** Approve for 1 year if the patient meets the following (A and B):
 - A)** Patient is ≥ 6 years of age; AND
 - B)** Patient meets ONE of the following (i, ii, iii, or iv):
 - i.** Patient meets BOTH of the following (a and b):
 - a)** Patient has tried oral iron supplementation; AND
 - b)** According to the prescriber, oral iron supplementation was ineffective or intolerable; OR
 - ii.** Patient has a condition which, per the prescriber, will interfere with oral iron absorption (e.g., inflammatory bowel disease, Crohn's disease); OR
 - iii.** Patient is currently receiving an erythroid stimulating agent; OR
Note: Examples of erythroid stimulating agents include an epoetin alfa product, a darbepoetin alfa product, or a methoxy polyethylene glycol-epoetin beta product.
 - iv.** The medication is being requested for cancer- or chemotherapy-related anemia.
- 4. Iron Deficiency Associated with Heart Failure.** Approve for 1 year if the patient meets the following (A and B):
 - A)** Patient is ≥ 6 years of age; AND
 - B)** Ferrlecit is being prescribed by or in consultation with a cardiologist or hematologist.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Ferrlecit is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Ferrlecit® [prescribing information]. Bridgewater, NJ: sanofi-aventis; March 2022.
2. Kidney Disease: Improving Global Outcomes (KDIGO) Anemia Work Group. KDIGO Clinical Practice Guideline for Anemia in Chronic Kidney Disease. *Kidney Int.* 2012;2(Suppl):279-335.
3. The NCCN Hematopoietic Growth Factors Guidelines in Oncology (version 2.2024 – December 12, 2023). © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on January 4, 2024.
4. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines [published correction appears in *J Am Coll Cardiol.* 2023 Apr 18;81(15):1551]. *J Am Coll Cardiol.* 2022;79(17):e263-e421.